

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>5</sup> :</b> <b>C12Q 1/00, G01N 33/53</b>	<b>A1</b>	<b>(11) International Publication Number:</b> • <b>WO 92/19761</b> <b>(43) International Publication Date:</b> 12 November 1992 (12.11.92)
<b>(21) International Application Number:</b> PCT/US92/03606 <b>(22) International Filing Date:</b> 1 May 1992 (01.05.92)  <b>(30) Priority data:</b> 695,805                      6 May 1991 (06.05.91)              US  <b>(71) Applicant:</b> THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 300 Lakeside Drive, 22nd Floor, Oakland, CA 94612 (US).  <b>(72) Inventors:</b> ROSEN, Steven ; 828 Clayton Street, San Francisco, CA 94117 (US). SINGER, Mark ; 1915 Grant Street, Berkeley, CA 94703 (US). IMAI, Yasuyuki ; 172 Locksley Avenue No.3, San Francisco, CA 94122 (US). YEDNOCK, Ted ; 46 Crest Road, Fairfax, CA 94930 (US).		<b>(74) Agents:</b> BASTIAN, Kevin, L. et al.; Townsend and Townsend, One Market Plaza, 2000 Steuart Tower, San Francisco, CA 94105 (US).  <b>(81) Designated States:</b> AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ASSAYS FOR INHIBITORS OF LEUKOCYTE ADHESION  <b>(57) Abstract</b> <p>The present invention provides novel assays for determining the ability of a test compound to inhibit intercellular adhesion mediated by a selectin receptor, such as the LHR. The assays involve contacting the test compound with the receptor and an isolated receptor-binding agent. The receptor-binding agent is a sulfated polysaccharide, a sulfated glycolipid, or a compound comprising the extracellular region of an endothelial cell surface glycoprotein. Also provided are compositions comprising a compound comprising the extracellular region of an endothelial cell surface protein, which is specifically recognized by lymphocyte homing receptors.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

ASSAYS FOR INHIBITORS OF LEUKOCYTE ADHESION

5           This invention was made with support under Grant (or Contract) No. NIH GM-23547, awarded by the DHHS. The U.S. Government has certain rights in this invention.

BACKGROUND OF THE INVENTION

10           The present invention relates to compositions and methods for identifying compounds useful in the inhibition of cellular adhesion involved in a number of pathological responses. In particular, the invention relates to assays for inhibitors of a class of receptors which mediate leukocyte extravasation and other responses.

15           Recent work has established that specialized cell surface receptors (termed here selectins) on endothelial cells and various circulating cells are involved in a number of intercellular interactions. For instance, an adhesion molecule on the surface of leukocytes, lymphocyte homing receptor (LHR),  
20           is known to be involved in the adhesive interactions of leukocytes with the endothelial lining of blood vessels. This adhesive interaction is a prerequisite for the movement of leukocytes from the blood to tissue sites where immune reactions and inflammatory reactions occur.

25           LHR (also known as gp90<sup>MEL</sup>, gp100<sup>MEL</sup>, gp110<sup>MEL</sup>, Mel-14 antigen, Leu8 antigen, TQ1 antigen, DREG antigen, LAM-1, selectin1, LECAM-1 and LEC-CAM-1, depending on animal species, leukocyte, and laboratory preference) is expressed on the surface of leukocytes, such as, lymphocytes, neutrophils,  
30           monocytes, and eosinophils (Gallatin, et al., Nature 303:30 (1983) and Lewinsohn, et al., J. Immunol. 138:4313 (1987)), LHR is known to mediate the adhesion of lymphocytes to specialized endothelial cells in lymph nodes, leading to the migration of blood-borne lymphocytes into the lymph node. On neutrophils  
35           and monocytes, it mediates the early interaction of these cells with endothelium of blood vessels at sites of inflammation.

          LHR is a lectin-like protein which performs its adhesive function by recognizing carbohydrate-containing

ligands on endothelial cells. Lectin-like receptors have also been found on endothelial cells and platelets. Endothelial leukocyte adhesion molecule-1 (ELAM-1) is present on endothelial cells and is involved in the recognition of various circulating cells by the endothelium. Granule membrane protein-140 (GMP-140) is present on the surface of platelets and endothelial cells, where it mediates platelet-leukocyte and endothelium-leukocyte interactions.

Recent work has established that these receptors share certain structural features. Each of the receptors in this class is a glycoprotein with a lectin-like domain, a region with homology to epidermal growth factor, and a region with homology to complement regulatory proteins (see, Springer, Nature, 346:425, 1989). The term "selectin" is used herein to refer to this class of lectin-like receptors.

There is currently an interest in developing highly specific competitive inhibitors of selectin-mediated cellular adhesion. Such inhibitors are useful in therapeutic regimens to treat various selectin-mediated disease responses. The inhibitors could also be used to target other pharmaceutical compounds, such as anti-inflammatory agents or anti-oxidants, to the sites of injury.

To date, however, insufficient understanding of the interaction of selectin receptors and their ligands has hindered these efforts. In addition, the prior art lacks rapid, economical methods for identifying inhibitors of selectin-mediated interactions. For example, in vitro intercellular adhesion assays have been used to test inhibition (see, e.g., Stamper and Woodruff, J. Exp. Med. 144:828-833 (1976)). These assays, however, are difficult to carry out and do not lend themselves to screening large numbers of test compounds. Also, comparisons of active compounds by quantitative dose-response studies is difficult using these assays. The present invention addresses these and related needs.

#### SUMMARY OF THE INVENTION

The present invention provides novel assays for determining the ability of a test compound to inhibit intercellular adhesion mediated by a selectin receptor, such as the LHR. The assays involve contacting the test compound with the receptor and an isolated receptor-binding agent. The receptor-binding agent is a phosphorylated polysaccharide, a sulfated polysaccharide, a sulfated glycolipid, or a compound comprising the extracellular region of an endothelial cell surface glycoprotein. The agent is preferably PPME, fucoidin, dextran sulfate, or a compound comprising the extracellular region of Sgp<sup>50</sup> or Sgp<sup>90</sup>.

The assays are typically carried out in a cell-free environment in which either the receptor or receptor-binding agent is immobilized on a solid surface. The assays may also be performed using intact cells and a fluorescence activated cell sorter. The ability of the test compound to inhibit binding between the receptor and the agent is detected in a number of ways, typically by using a labeled receptor binding agent.

Also disclosed are compositions comprising the extracellular region of an endothelial cell surface glycoprotein. The extracellular region comprises a carbohydrate ligand specifically recognized by LHR. The glycoprotein is typically Sgp<sup>50</sup> or Sgp<sup>90</sup>.

Finally, compositions comprising inhibitor compounds identified by the above methods are disclosed. The inhibitors can be used in pharmaceutical compositions to treat various selectin-mediated disease states.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides novel assays useful in identifying inhibitors of selectin-mediated intercellular adhesion. Also provided are isolated endothelial cell surface glycoproteins which selectively bind lymphocyte homing receptors. Compounds comprising the extracellular region of these glycoproteins are particularly useful in the assays of the present invention.

As discussed above, selectins, also known as the "LEC-CAM" family of cell adhesion molecules, are unique cell surface glycoproteins. These receptors are involved in a variety of intercellular interactions. For instance, the trafficking of lymphocytes from the blood into secondary lymphoid organs, such as lymph nodes and gut-associated Peyer's patches, is known to be initiated by an adhesive interaction between specialized endothelial cells of high endothelial venules (HEV) and LHRs on lymphocytes. Berg et al., Immunol. Rev. 108:5-18 (1989); Duijvestijn and Hamann, Immunol. Today 10:23-28 (1989); Woodruff et al., Ann. Rev. Immunol. 5:201-222 (1987); Yednock and Rosen, Adv. Immunol. 54:313-378 (1989); Stoolman, Cell 56:907-910 (1989); Gallatin et al., Cell 44:673-680 (1986); Rosen, Curr. Opin. Cell. Biol. 1:913-919 (1989).

The endothelial ligands recognized by LHR molecules are postulated to be distinctive for the different lymphoid organs and as such are proposed to be responsible for regulating the lymphocyte populations that enter each class of lymphoid organ. Strong support for organ-specific HEV determinants has come with the discovery of the mouse "vascular addressin" antigens, defined by the panel of MECA monoclonal antibodies (Streeter et al., Nature 331:41-46 (1988) and Streeter et al., J. Cell. Biol. 107:1853-1962 (1988)).

The lectin domain on LHR for lymph nodes in humans and mice was initially inferred based upon the ability of specific phosphorylated monosaccharides, such as mannose-6-phosphate (M6P), and specific polysaccharides to prevent lymphocyte attachment to HEV (Stoolman and Rosen, J. Cell Biol. 96:722-729 (1983); Stoolman et al., J. Cell Biol. 99:1535-1540 (1984); Yednock et al., J. Cell Biol. 104:713-723 (1987); Stoolman et al., Blood 70:1842-1850 (1987); Stoolman and Ebling, J. Clin. Invest. 84:1196-1205 (1989)). Notable among the active polysaccharides are PPME (a phosphate-rich mannan core) and fucoidin (a sulfated, fucose-rich polymer). This carbohydrate-binding activity depends on the presence of calcium, which is also required for the attachment of lymphocytes to HEV.

From the lectin nature of LHR, the ligands on lymph node HEV are presumed to bear a carbohydrate-based recognition determinant. Early studies demonstrated that the adhesive sites on peripheral lymph node HEV are periodate sensitive (Rosen et al., Science 228:1005-1007 (1985)) indicating a requirement for carbohydrate. Subsequently, it was demonstrated that sialidase treatment of HEV, in vitro or in vivo, selectively eliminates lymphocyte attachment to peripheral lymph node HEV but has no effect on the binding to Peyer's patch HEV (Rosen et al., J. Immunol. 142:1895-1902 (1989)). In addition, exposure of peripheral lymph node tissue sections to Limax flavus agglutinin, a sialic acid-specific lectin, prevents lymphocyte attachment to HEV (True et al., J. Cell Biol. 111:2757-2764 (1990)).

The biochemical nature of the ligands, however, has not been defined. Adhesion-blocking activity and selective staining of peripheral lymph node HEV have been shown by MECA-79, a monoclonal antibody which apparently recognizes a complex of HEV cell surface proteins (EPO Publication No. 0303463 and Butcher, Am. J. Pathol. 136:3-12 (1990)). The present invention provides the first evidence that LHR specifically binds a surface glycoprotein recognized by this antibody. In particular evidence provided here establishes that LHR recognizes a sulfated, fucosylated and sialylated glycoprotein of about 50 kd. A glycoprotein of 90 kd with similar characteristics is also identified.

Other selectins have also been extensively studied. ELAM-1 is inducibly expressed on vascular endothelial cells (Bevilacqua et al., Science 243:1160 (1989) and Hession et al., Proc. Nat'l. Acad. Sci., 87:1673-1677 (1990)). This receptor has been demonstrated to be induced by inflammatory cytokines such as interleukin I $\beta$  (IL-I $\beta$ ) and tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), as well as bacterial endotoxin (lipopolysaccharide) (see, Bevilacqua et al., Proc. Natl. Acad. Sci., 84:9238-9242 (1987)). These compounds act directly on endothelial cells in vitro to substantially augment polymorphonuclear leukocyte (neutrophil), and monocyte adhesion (Bevilacqua et al., Proc. Natl. Acad. Sci., supra). Detailed structures have been

proposed for an oligosaccharide moiety recognized by ELAM-1 (Philips et al. Science 250:1130-1132 (1990) and Walz et al. Science 250:1132-1135 (1990)).

5 A third member of the selectin family, GMP-140, is a membrane glycoprotein of platelet and endothelial secretory granules (Geng et al., Nature, 343, 757-760 (1990)). Activated platelets which express GMP-140 on their surface are known to bind to monocytes and neutrophils (Jungi et al., Blood 67:629-636 (1986)), and also to monocyte-like cell lines, e.g., HL60  
10 and U937 (Jungi et al., supra; Silverstein et al., J. Clin. Invest. 79:867-874 (1987)). GMP-140 is an alpha granule membrane protein of molecular weight 140,000 that is expressed on the surface of activated platelets upon platelet stimulation and granule secretion (Hsu-Lin et al., J. Biol. Chem. 259:9121-9126 (1984); Stenberg et al., J. Cell Biol. 101:880-886 (1985);  
15 Berman et al., J. Clin. Invest. 78:130-137 (1986)). It is also found in megakaryocytes (Beckstead et al., Blood 67:285-293 (1986)), and in endothelial cells (McEver et al., Blood 70:355a (1987)) within the Weibel-Palade bodies (Bonfanti et al., Blood  
20 73:1109-1112 (1989)). Furie et al. U.S. Patent No. 4,783,330, describe monoclonal antibodies reactive with GPM-140.

The structure and function of selectin receptors has been elucidated by cloning and expression of full length cDNA encoding each of the above receptors (see, e.g., Bevilacqua et  
25 al., Science, supra, (ELAM-1), Geng et al., supra, (GMP 140), and Lasky et al., Cell 56:1045-1055 (1989) (LHR). The extracellular portion of selectins can be divided into three segments based on homologies to previously described proteins. The N-terminal region (about 120 amino acids) is related to the  
30 C-type mammalian lectin protein family as described by Drickamer, J. Biol. Chem., 263: 9557-9560 (1988) that includes low affinity IgE receptor CD23. A polypeptide segment follows, which has a sequence that is related to proteins containing the epidermal growth factor (EGF) motif. Lastly, after the EGF  
35 domain are one or more tandem repetitive motifs of about 60 amino acids each, related to those found in a family of complement regulatory proteins.



In one aspect, the present invention is directed to methods for assaying test compounds for the ability to inhibit selectin-mediated responses. Ideally, the assays of the present invention allow large scale in vitro screening of a variety of compounds.

Inhibition of a number of binding interactions can be assayed. For instance, inhibition of the binding between a monoclonal antibody and an isolated selectin receptor can be used. Numerous types of competitive assays involving immunoglobulins are known (see, e.g., U.S. Patents No. 3,376,110, 4,016,043, and Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Publications, N.Y. (1988)).

The in vitro assays of the present invention are typically assays which detect the ability of a test compound to inhibit binding between the receptor and an isolated receptor-binding agent. The term "isolated receptor-binding agent" is used to refer to an agent that is not associated with an intact cell membrane. Thus, intact cells which express ligand molecules recognized by selectin receptors are not used in the assays of the present invention. The receptor-binding agents are typically prepared from appropriate cells according to standard methods, purchased in a purified state, or synthetically produced. Although the agents are not associated with intact cell membranes they may be associated with cell membrane fragments or incorporated into artificial lipid membranes (e.g. liposomes).

Unlike prior art methods, the present assays are not based on measuring the presence or absence of intercellular adhesion but are typically carried out in a cell-free environment. The term "a cell-free environment" is used here to indicate that intact cells are substantially absent from the assay. The assay is considered to be cell-free despite the presence of cellular debris or cell membrane fragments which comprise a component of the assay. In some formats, however, (e.g. those using a fluorescence activated cell sorter) intact cells, such as lymphocytes, which express the receptor of interest are used.

A number of compounds known to block intercellular adhesion can be used as the receptor-binding agent in the present invention. For instance, phosphorylated monosaccharides, such as mannose-6-phosphate and fructose-1-phosphate inhibit lymphocyte attachment to HEV in in vitro cellular assays. Polysaccharides and glycolipids, have also been shown to inhibit in vitro binding of these cells (Stoolman et al., supra; Yednock et al., supra, and Yednock et al., J. Cell Biol., 104:725-731 (1987)).

The polysaccharides of the present invention are phosphorylated or sulfated. Phosphorylated polysaccharides include the phosphomannan monoester core from Hansenula hostii (PPME). Sulfated polysaccharides include fucoidin and dextran sulfate. Sulfated glycolipids of the invention include sulfatide. Using the novel assays claimed here, however, one of skill can easily identify other polysaccharides and glycolipids within the scope of the claims which are effective receptor-binding agents. For instance, Example 1, below, provides a method suitable for assaying such compounds. The results provided there indicate that different compounds have different inhibitory activities.

Receptor-binding agents of the present invention can be easily prepared from commonly available starting materials. Polysaccharides and glycolipids are isolated from animal, plant, fungal or prokaryotic cells according to standard techniques. For instance, PPME is purified from crude yeast mannan by the method of Slodki et al., Biochim. Biophys. Acta, 304:449-456 (1973). Briefly, the phosphomannan is acid hydrolyzed. After neutralization, the phosphomannan core is precipitated and rehydrated in water. Contaminating protein is removed by water:chloroform:butanol extraction. The compounds can also be chemically synthesized according to standard techniques (see, e.g., Khadem, Carbohydrate Chemistry (Academic Press, San Diego, CA, 1988)). Alternatively, many polysaccharides (such as fucoidin) and glycolipids can be purchased from chemical supply companies, such as Sigma Chemical Co. (St. Louis, MO) and Aldrich Chemical Co. (Milwaukee, WI).

The selectin receptor used in the assays of the present invention may or may not be associated with an intact cell. Typically, the receptor is purified from its native environment before use in the assay. As discussed above, cDNA encoding each of the selectin receptors has been isolated. Thus, the receptors can be recombinantly produced using standard methods well known to those skilled in the art. For a review of standard molecular biological techniques see Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed. (Cold Spring Harbor Press, N.Y., 1989). In addition, using standard recombinant DNA techniques, mutations can be induced to obtain proteins with altered amino acid sequences. Typically, substitutions, deletions or additions are introduced which provide desired characteristics. For instance, increased solubility can be achieved by elimination of the hydrophobic transmembrane region of the protein. In addition, soluble chimeric receptors comprising the constant region of an immunoglobulin molecule can also be produced (Watson et al., J. Cell Biol. 110:2221-2229 (1990), and Watson et al., Nature 349:164-167 (1991).

The test compound to be screened will usually be a synthetic or naturally-produced biomolecule, such as a carbohydrate (e.g., oligosaccharide) or glycoconjugate. It can also be a peptide, polypeptide, protein (e.g., monoclonal antibody), nucleic acid, and the like. The test compound is typically a relatively small molecule with a molecular weight less than about 10 kD, preferably less than about 5 kD. The compounds are synthetically produced using standard methods for synthesizing oligosaccharides (Khadem, supra). Methods for synthesizing polypeptides of defined composition are well known in the art (see, Atherton et al. Solid Phase Peptide Synthesis (IRL Press, Oxford, 1989)). If the synthetic test compounds are polymeric (e.g., polypeptides or polysaccharides) they are preferably altered in a systematic way to identify the sequence of monomers which have the desired effect (see, e.g., U.S. Patent No. 4,833,092). Test compounds may also be isolated from any natural source, such as animal, plant, fungal, or prokaryotic cells in accordance with standard procedures.

The assays of the present invention are particularly useful in identifying compounds which act as antagonists of a ligand molecule. Antagonists are compounds which reverse the physiological effect of a ligand or exclude binding of the ligand to the receptor. An antagonist competes directly or indirectly with the ligand for the receptor binding site and, thus, reduces the proportion of ligand molecules bound to the receptor. Typically, an antagonist will be the topographical equivalent of the natural ligand and will compete directly with the ligand for the binding site on the selectin. Such a compound is referred to here as a "mimetic." A ligand mimetic is a molecule that conformationally and functionally serves as substitute for the natural ligand recognized by a selectin receptor. Alternatively, if the ligand and the test compound can bind the receptor simultaneously, the compound may act non-competitively. A non-competitive inhibitor acts by decreasing or inhibiting the subsequent physiological effects of receptor-ligand interactions rather than by diminishing the proportion of ligand molecules bound to the receptor.

The assays of the present invention can also be used to identify synthetic or naturally occurring agonists. Agonists are compounds which bind the receptor and initiate a physiological response similar to that of the natural ligand.

The inhibitors of selectin-ligand interaction identified by the assays of the present invention are useful in treating a number of selectin-mediated disease responses. For instance, selectins play an important role in recruitment of leukocytes to the sites of injury, particularly inflammation. The inhibitors therefore may be administered locally or systemically to control tissue damage associated with such injuries. Moreover, because of the specificity of such inhibitors for sites of inflammation, these compositions will be more effective and less likely to cause complications when compared to traditional anti-inflammatory agents.

Pharmaceutical compositions comprising the inhibitors can be used to block or inhibit cellular adhesion associated with a number of disorders. For instance, a number of inflammatory disorders are associated with selectins expressed

on vascular endothelial cells and platelets. The term "inflammation" is used here to refer to reactions of both the specific and non-specific defense systems. A specific defense system reaction is a specific immune system reaction to an antigen. Examples of specific defense system reactions include antibody response to antigens, such as viruses, and delayed-type hypersensitivity. A non-specific defense system reaction is an inflammatory response mediated by leukocytes generally incapable of immunological memory. Such cells include macrophages, eosinophils and neutrophils. Examples of non-specific reactions include the immediate swelling after a bee sting, and the collection of PMN leukocytes at sites of bacterial infection (e.g., pulmonary infiltrates in bacterial pneumonias and pus formation in abscesses).

Other treatable disorders include, e.g., rheumatoid arthritis, post-ischemic leukocyte-mediated tissue damage (reperfusion injury), frost-bite injury or shock, acute leukocyte-mediated lung injury (e.g., adult respiratory distress syndrome), asthma, traumatic shock, septic shock, and acute and chronic inflammation, including atopic dermatitis, psoriasis, and inflammatory bowel disease. Various platelet-mediated pathologies such as atherosclerosis and clotting can also be treated. In addition, tumor metastasis can be inhibited or prevented by inhibiting the adhesion of circulating cancer cells. Examples include carcinoma of the colon and melanoma.

Thus, the present invention also provides pharmaceutical compositions which can be used in treating the aforementioned conditions. The pharmaceutical compositions are comprised of inhibitory compounds together with pharmaceutically acceptable carriers. The pharmaceutical compositions can be prepared according to standard methods (see Remington's Pharmaceutical Sciences, Mack Publishing Co., Philadelphia, PA, 19th ed. (1985) which is incorporated herein by reference). The pharmaceutical compositions are suitable for use in a variety of drug delivery systems. For a brief review of present methods for drug delivery, see, Langer, Science 249:1527-1533 (1990).

In one embodiment, the inhibitors can be used to target conventional anti-inflammatory drugs or other agents to specific sites of tissue injury. By using a selectin-binding moiety to target a drug to a selectin receptor on, e.g., a vascular endothelial cell, such drugs can achieve higher concentrations at sites of injury. Side effects from the conventional anti-inflammatory agents can be substantially alleviated by the lower dosages, the localization of the agent at the injury sites and/or the encapsulation of the agent prior to delivery. Targeting can be achieved by directly or indirectly linking the inhibitor to the anti-inflammatory agent. For instance, liposomes filled with the anti-inflammatory agent can be constructed which incorporate the inhibitor in the lipid membrane (see, Langer, supra). When the liposomes are brought into proximity of the affected cells, they deliver the elected therapeutic compositions.

The pharmaceutical compositions containing the inhibitors can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, compositions are administered to a patient already suffering from a disease, as described above, in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will, of course, depend on the severity of the disease and the weight and general state of the patient.

In prophylactic applications, compositions containing the inhibitors are administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight.

The present invention also provides lymph node endothelial cell surface sulfated glycoproteins which comprise oligosaccharide biological ligands specifically recognized by LHR for lymph nodes. As described more fully below, two such glycoproteins, Sgp<sup>50</sup> and Sgp<sup>90</sup>, have been identified. Having identified ligand-bearing glycoproteins, one of skill will

recognize that a number of modifications of the glycoproteins that do not significantly alter the LHR binding activity are possible. Such modifications include enzymatic or chemical treatment of the proteins to produce fragments that comprise the carbohydrate ligand recognized by LHR. For instance, fragments of the extracellular region of the proteins can be obtained by treatment of the isolated glycoproteins with an appropriate protease such as trypsin, pronase, papain, pepsin and the like.

The extracellular region of the cell surface glycoproteins includes all sequences from the proteins outside the transmembrane and intracellular regions. The extracellular region of the glycoproteins of the present invention comprises a carbohydrate ligand specifically recognized by LHR. The extracellular region may also contain sequences from the transmembrane region (less than about 10 amino acids), so long as solubility is not substantially affected. The term "compound comprising the extracellular region" includes any compound in which the extracellular region, or fragment thereof, is conjugated to a second moiety. The term also embraces the isolated extracellular region and the isolated full length glycoprotein. An "isolated compound comprising the extracellular region" includes such a compound (e.g., a full length glycoprotein) in other than its native state, that is, not associated with an endothelial cell. For instance, the compound may be recombinantly produced, solubilized from the appropriate cell, or associated with a liposome.

Analysis of sulfated the glycoproteins of the present invention has revealed that the oligosaccharide moieties recognized by LHR are O-linked. Thus, they can be cleaved from the protein backbones by beta elimination and borohydride reduction according to standard techniques (see, e.g., Fukuda, Meth. Enzymol. 179:17-29 (1989)).

The sulfated glycoproteins, or fragments thereof, can be isolated using soluble LHR as described below. The isolated soluble molecules are then used in the assays of the present invention. These molecules are particularly useful in the invention because inhibition of the interaction of the LHR with

the actual biological ligand is assayed. The glycoproteins are also suitable as inhibitors of LHR-mediated cellular adhesion in the pharmaceutical compositions, described above. In either case, the glycoproteins can be used as they are isolated or they can be conjugated to a variety of other compounds to confer any number of desired characteristics, such as improved solubility, serum half-life and the like. Alternatively, neoglycoproteins or neoglycolipids can be prepared based on the carbohydrate chains of the glycoproteins using methods well known in the art (see, e.g., Stowell et al., Adv. Carb. Chem and Biochem. 37:225-281 (1980) and Childs et al. Biochem. J., 262:131-138 (1989)).

In the assays of the present invention, any component of the assay, typically the receptor, is bound to a solid surface. For instance, in the case of LHR-based assays, purified LHR is coated on a solid surface and the ability of the test compound to inhibit binding between the receptor and a LHR-binding agent is detected.

Many methods for immobilizing biomolecules on solid surfaces are known in the art. For instance, the solid surface may be a membrane (e.g., nitrocellulose), a microtiter dish (e.g., PVC or polystyrene) or a bead. The desired component may be covalently bound or noncovalently attached through unspecific bonding.

A wide variety of organic and inorganic polymers, both natural and synthetic may be employed as the material for the solid surface. Illustrative polymers include polyethylene, polypropylene, poly(4-methylbutene), polystyrene, polymethacrylate, poly(ethylene terephthalate), rayon, nylon, poly(vinyl butyrate), silicones, polyformaldehyde, cellulose, cellulose acetate, nitrocellulose, etc. Other materials which may be employed, include paper, glasses, ceramics, metals, metalloids, semiconductive materials, cermets or the like. In addition are included substances that form gels, such as proteins, e.g., gelatins, lipopolysaccharides, silicates, agarose and polyacrylamides or polymers which form several aqueous phases, such as dextrans, polyalkylene glycols (alkylene of 2 to 3 carbon atoms) or surfactants e.g.



amphophilic compounds, such as phospholipids, long chain (12-24 carbon atoms) alkyl ammonium salts and the like. Where the solid surface is porous, various pore sizes may be employed depending upon the nature of the system.

5 In preparing the surface, a plurality of different materials may be employed, particularly as laminates, to obtain various properties. For example, protein coatings, such as gelatin can be employed to avoid non-specific binding, simplify covalent conjugation, enhance signal detection or the like.

10 If covalent bonding between a compound and the surface is desired, the surface will usually be polyfunctional or be capable of being polyfunctionalized. Functional groups which may be present on the surface and used for linking can include carboxylic acids, aldehydes, amino groups, cyano  
15 groups, ethylenic groups, hydroxyl groups, mercapto groups and the like. The manner of linking a wide variety of compounds to various surfaces is well known and is amply illustrated in the literature. See for example Chibata, Immobilized Enzymes, Chibata, Halsted Press, New York, 1978, and Cuatrecasas, J. Biol. Chem. 245:3059 (1970).

20 In addition to covalent bonding, various methods for noncovalently binding an assay component can be used. Noncovalent binding is typically nonspecific absorption of a compound to the surface. Typically, the surface is blocked  
25 with a second compound to prevent nonspecific binding of labeled assay components. Alternatively, the surface is designed such that it nonspecifically binds one component but does not significantly bind another. For example, a surface bearing a lectin such as Concanavalin A will bind a  
30 carbohydrate containing compound but not a labeled protein that lacks glycosylation. Various solid surfaces for use in noncovalent attachment of assay components are reviewed in U.S. Patent Nos. 4,447,576 and 4,254,082.

35 Many assay formats employ labeled assay components, typically the selectin-binding agent. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide variety of labels may be used. The component may be labeled by any one of

several methods. A common method of detection is the use of autoradiography with  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^{32}\text{P}$  labeled compounds or the like. The choice of radioactive isotope depends on research preferences due to ease of synthesis, varying  
5 stability, and half lives of the selected isotopes. Other non-radioactive labels include ligands which bind to labeled antibodies, fluorophores, chemiluminescent agents, enzymes, and antibodies which can serve as specific binding pair members for a labeled ligand. The choice of label depends on sensitivity  
10 required, ease of conjugation with the compound, stability requirements, and available instrumentation.

Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand then binds to an  
15 anti-ligand (e.g., streptavidin) molecule which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. Ligands and anti-ligands may be varied widely. Where a ligand has a natural anti-ligand, for  
20 example, biotin, thyroxine, and cortisol, it can be used in conjunction with the labeled, naturally occurring anti-ligands. Alternatively, any haptenic or antigenic compound can be used in combination with an antibody.

The molecules can also be conjugated directly to  
25 signal generating compounds, e.g., by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidoreductases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives,  
30 rhodamine and its derivatives, dansyl, umbelliferone, etc. Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g., luminol. For a review of various signal producing systems which may be used, see, U.S. Patent No. 4,391,904.

35 In other embodiments, a fluorescence-activated cell sorter (FACS) is used to isolate selectin bearing cells and quantify inhibitory effects of test compounds. Typically, receptor-binding agents are labeled with a fluorescent compound

and the amount of labeled agent bound to the cell surface is determined by flow cytometry. In the FACS, a stream of cells flows past a laser beam which causes the labeled cells to emit light. The stream is broken into droplets, each containing a single cell. Droplets containing a fluorescent cell are given a charge, which allows them to be separated from droplets containing no cell or an unlabeled cell. In addition, to sorting the apparatus can be used to measure the level of fluorescence in labeled cells. For a general discussion of fluorescence activated cell sorting and flow cytometry see Parks et al., in Fundamental Immunology, Paul, ed. (Raven Press, 2d. ed. 1989).

In the assays of the present invention, fluorescent-labeled agents (such as PPME and fucoidin) are used to detect selectin receptors on cells such as leukocytes, platelets, and endothelial cells. For instance, flow cytometry can be used to detect LHR on viable lymphocytes and inhibitors can be easily determined. In addition, a quantitative measure of a receptor's activity on a cell surface can be determined.

The following examples are offered by way of illustration, not by way of limitation.

#### Example 1

##### PPME-based ELISA Assay with Anti-PPME Antibody

This example shows the ability of assays of the present invention to identify inhibitors of LHR-mediated intercellular adhesion. The illustrated assay uses LHR coated on the wells of microtiter plates. Binding of PPME to the coated wells is detected with a rabbit antibody to PPME.

To perform the assay, LHR for lymph nodes purified by immunoaffinity chromatography as previously described (Lasky et al. supra; Geoffroy and Rosen, J. Cell Biol. 109:2463-2469 (1989)) was equilibrated in Dulbecco's PBS (0.68 mM calcium, 0.49 mM magnesium) containing either 1 mM CHAPS or 40 mM octyl $\beta$ -D-glucopyranoside. The PPME-binding activity of LHR appeared to be more stable in octyl $\beta$ -D-glucopyranoside than in CHAPS.

10  $\mu$ l of LHR solution at specified concentrations was diluted into 90  $\mu$ l of PBS in the wells of Immulon-2 microtiter plates (Dynatech Laboratories, Inc., Chantilly, VA). The dilution factor was chosen to reduce the detergent

5 concentration below the critical micellar concentration.

Adsorption was allowed to proceed overnight at 4°C. The wells were washed three times with PBS containing 0.1% Tween 20 (PBS-Tween). 200  $\mu$ l of PBS containing 3% BSA and 0.02% NaN<sub>3</sub> was then added to each well to block nonspecific binding sites.

10 After a 2-h incubation at 22°C with agitation, the wells were washed and then received 100  $\mu$ l of PPME in 3% BSA-PBS or 3% BSA-PBS alone (control for background). After a 1-h incubation at 4°C, the wells were washed to remove unbound polysaccharide and 100  $\mu$ l of rabbit anti-PPME antiserum (1:100  
15 dilution into 3% BSA in PBS-Tween) was added to each well.

Anti-PPME antiserum was prepared by injection New England white rabbits with 100  $\mu$ g of PPME emulsified in 2 ml Freund's complete adjuvant (Difco Laboratories, Detroit, MI). The antigen was administered in multiple intradermal and  
20 subcutaneous sites.

The incubation of the test compound in the presence of LHR and PPME and all subsequent incubations were allowed to proceed at 22°C with agitation. After 1 h of exposure to anti-PPME, the wells were washed and 100  $\mu$ l per well of biotinylated  
25 goat anti-rabbit IgG (1:200 dilution into 3% BSA in PBS-Tween) was added and allowed to react for 1 h. The wells were again washed before the addition of 50  $\mu$ l per well of Vector ABC-AP reagent in PBS-Tween made according to the manufacturer's specifications. After 30 min., the wells were washed again and  
30 color development was initiated by the addition of 100  $\mu$ l per well of substrate solution (p-nitrophenylphosphate, 1 mg/ml in 10% [vol/vol] diethanolamine-HCl buffer, pH 9.8).

Optical density readings were made at 405 nm with a microplate reader (BioRad Laboratories, Model 3550; Richmond,  
35 CA). Mean values are determined for two to three replicate wells. With PPME added in excess, a saturation level of binding was observed at  $\geq 0.4$   $\mu$ g of LHR per well.

In assays measuring the ability of test components to compete with PPME for binding to immobilized LHR, a concentration of 0.2-0.5  $\mu\text{g}$  per well of LHR was added and the substances (diluted into 50  $\mu\text{l}$  of PBS) were allowed to react with the antigen for 1 h at 4°C. PPME (50  $\mu\text{l}$ , in 6% BSA-PBS) was added to each well, and the plate was incubated for an additional 1 h at 4°C. The plate was then processed as described above.

The concentrations of the test compounds are expressed as final concentrations after addition of the PPME. To test lipids as inhibitors, samples dissolved in chloroform-methanol 1:1 were dried in glass test tubes under a stream of nitrogen, and then suspended in PBS by brief sonication (model D-50; Branson Sonic Power Co., Chicago IL). All of the inhibitory effects documented in this example were observed in two independent experiments. The ability of the assay to detect the inhibitory effects of various test compounds is presented in Tables 1 and 2.

Table 1

## Inhibitory Activities of Sulfated Macromolecules

25	Substance	Concentration for 50% inhibition
30		$\mu\text{g/ml}$
	sulfatide (Matreya, Inc.)	ND
	sulfatide (Sigma Chemical Co.)	0.7-1.3
	cholesterol-3-sulfate	>100
	chondroitin sulfate A	>1,000
35	chondroitin sulfate B	>1,000
	chondroitin sulfate C	>1,000
	keratan sulfate	>1,000
	fucoidin (Sigma Chemical Co.)	ND
	heparan sulfate	>1,000
40	fucoidin (CN K&K Laboratories)	0.1-0.4
	heparin (porcine intestinal)	100
	heparin (bovine lung)	19
	heparin (porcine mucosa)	110
	egg jelly fucan	0.2
45	polyvinylsulfate	>1,000

Table 2

## Effects of Lipids on PPME Binding to LHR

5	Experiment	Inhibitor	$\mu\text{g/ml}$	OD 405 nm
10				
	1	None	-	0.596 $\pm$ 0.025
		Sulfatide (Sigma Chemical Co.)	10	0.030 $\pm$ 0.005
		Sulfatide (Metrya, Inc.)	10	0.019 $\pm$ 0.006
15		Galactosylceramide	10	0.668 $\pm$ 0.021
		Gal(6-SO <sub>4</sub> ) $\beta$ 1-1 Cer	10	0.612 $\pm$ 0.016
		Cholesterol 3-SO <sub>4</sub>	10	0.583 $\pm$ 0.014
20	2	None	-	0.291 $\pm$ 0.065
		Sulfatide (Sigma Chemical Co.)	10	0.023 $\pm$ 0.018
		G <sub>M1</sub>	10	0.208 $\pm$ 0.014
		G <sub>D1a</sub>	10	0.328 $\pm$ 0.017
		G <sub>D1b</sub>	10	0.256 $\pm$ 0.053
		G <sub>T1b</sub>	10	0.287 $\pm$ 0.017
25	3	None	-	0.362 $\pm$ 0.064
		Sulfatide (Metrya, Inc.)	100	0.0
		G <sub>M1</sub>	100	0.301 $\pm$ 0.026
		G <sub>M2</sub>	100	0.248 $\pm$ 0.042
30		G <sub>M3</sub>	100	0.378 $\pm$ 0.019
	4	None	-	0.431 $\pm$ 0.047
		Sulfatide (Metrya, Inc.)	10	0.0
		G <sub>D1a</sub>	100	0.331 $\pm$ 0.025
35		G <sub>D1b</sub>	100	0.313 $\pm$ 0.011
		G <sub>T1b</sub>	100	0.257 $\pm$ 0.026
		G <sub>D3</sub>	100	0.555 $\pm$ 0.021
		G <sub>Q1b</sub>	100	0.288 $\pm$ 0.032
40				

Structures of the Glycolipids tested in Table 2 are as follows: Galactosylceramide, Gal $\beta$ 1-1Cer; Sulfatide, Gal(3-SO<sub>4</sub>) $\beta$ 1-1Cer; G<sub>M1</sub>, Gal $\beta$ 1-3GalNAc $\beta$ 1-4[Neu5Ac $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>D1a</sub>, NeuAc $\alpha$ 2,3Gal $\beta$ 1-3GalNAc $\beta$ 11-4[NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>D1b</sub>, Gal $\beta$ 1-3GalNAc $\beta$ 1-4[NeuAc $\alpha$ 2-8NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>T1b</sub>, NeuAc $\alpha$ 2-3GalNAc $\beta$ 1-4[NeuAc $\alpha$ 2-8NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>M2</sub>, GalNAc $\beta$ 11-4[NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>M3</sub> NeuAc $\alpha$ 2-3Gal $\beta$ 1-4Glc $\beta$ 1-1Cer. G<sub>D3</sub>, NeuAc $\alpha$ 2-8NeuAc $\alpha$ 2-3Gal $\beta$ 1-4Glc $\beta$ 1-1Cer; G<sub>Q1b</sub>, NeuAc $\alpha$ 2-8NeuAc $\alpha$ 2-3Gal $\beta$ 1-3GalNAc $\beta$ 1-4[NeuAc $\alpha$ 2-8NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc $\beta$ 1-1Cer. Three replicates were performed with each glycolipid.

Example 2ELISA Assay with fluoresceinated-PPME or  
fluoresceinated-fucoidin

In this assay, native or recombinant LHR was coated  
5 onto plastic wells of an ELISA plate as in assay 1 above. A  
fluorescein conjugate of either PPME (i.e., fl-PPME) or of  
fucoidin (fl-fucoidin) was allowed to bind to the immobilized  
receptor. The bound conjugate was detected with an alkaline  
phosphatase-labeled, affinity-purified rabbit antibody to  
10 fluorescein (Biodesign, Inc., Kennebunkport, ME). The bound  
antibody was detected with a standard colorimetric assay for  
alkaline phosphatase in a microplate reader as described above.  
Binding of the fluorescein conjugates to the receptor  
approaches saturation and was substantially (in the case of fl-  
15 fucoidin) or completed inhibited (in the case of fl-PPME) by  
calcium chelation or by MEL-14.

The fluorescein conjugates were prepared by the  
procedure of Glabe et al., Anal. Biochem., 130:287-294 (1983).  
Briefly, the polysaccharides were activated by cyanogen bromide  
20 treatment and the reacted with fluoresceinamine. Conjugates  
are then separated from free fluoresceinamine by gel filtration  
according to standard procedures.

Example 3ELISA Assay with Sulfatide

In this example, the glycolipid sulfatide, was coated  
onto the wells of a polystyrene microtiter plate by standard  
absorption procedures. A recombinant LHR conjugate, LHR-IgG (a  
soluble molecule engineered with a human immunoglobulin Fc  
30 region), was allowed to bind to the immobilized sulfatide. The  
LHR-IgG based upon the LHR for lymph nodes was prepared as  
described in Watson et al. J. Cell Biol. 110:2221-2229 (1990).

The bound recombinant receptor was detected by  
sequential application of a biotinylated protein A (which  
35 reacts with the Fc region) and the alkaline phosphatase-ABC  
system. The reaction product was measured, as described above,  
with the microplate ELISA reader. A major component of the

signal was inhibited by EGTA (a calcium chelator) or by the MEL-14 antibody.

#### Example 4

##### Flow Cytometry Assay with fl-PPME or fl-fucoidin

The fluorescein conjugates of PPME and fucoidin (i.e., fl-PPME and fl-fucoidin), described in example 2 above, were used to detect LHR on the surface of viable lymphocytes by flow cytometry. Lymphocytes were exposed to the conjugates, washed, and the amount of conjugate bound to the cell surface was directly determined with a FACSCAN flow cytometer. The majority of lymphocytes reacted with each of the conjugates. The signals were substantially inhibited by EGTA (a calcium chelator) or by the MEL-14 antibody.



## Example 5

Identification of Surface Glycoproteins on  
Endothelial Cells Recognized by LHR

5           This example shows that recombinant LHR selectively binds  $^{35}\text{SO}_4$  labeled macromolecules from lymph nodes. In particular, two sulfated, fucosylated and sialylated glycoproteins have been identified.

10   A.   Metabolic Labeling of Organs with  $^{35}\text{S}$ -sulfate

          Mesenteric or peripheral (cervical, brachial, axillary) lymph nodes were collected from 8-16 week old female ICR mice. The lymph nodes were cut into 1 mm thick slices with a razor blade and the slices (typically, 0.2 g of wet weight) were suspended in 1 ml of RPMI-1640 containing 25 mM HEPES, 100 U/ml Penicillin G, 100  $\mu\text{g/ml}$  streptomycin, and 200  $\mu\text{Ci}$  carrier-free [ $^{35}\text{S}$ ] sodium sulfate (ICN Biochemicals Inc., Costa Mesa, CA) according to the procedure of Ager, J. Cell Sci., 87:133-144 (1987). After incubation at 37°C for 4 hr, the slices were washed extensively in Dulbecco's phosphate-buffered saline (PBS), and then homogenized in 1 ml of lysis buffer (2% Triton X-100 in PBS containing 1 mM PMSF, 1% (v/v) aprotinin, 10  $\mu\text{g/ml}$  pepstatin, 0.02%  $\text{NaN}_3$ ) with a Potter-Elvehjem homogenizer on ice. Lysis was continued for 1 hr on a rocker at 4°C. The lysate was centrifuged at 10,000 x g for 1 hr at 4°C. EDTA was added to the supernatant at a final concentration of 2 mM and the supernatant was precleared by rocking with Affi-Gel Protein A (250  $\mu\text{l}$  of packed beads, BioRad Laboratories, Richmond, CA) overnight at 4°C.

30

B.   Identification of the Components Adsorbed to LHR-IgG Beads

          Affi-Gel Protein A (10  $\mu\text{l}$  packed beads) was incubated with 30  $\mu\text{g}$  of either LHR-IgG (prepared as described above), CD4-IgG (prepared according to Capon et al., Nature 337:525-531 (1989), or human IgG, (Calbiochem, La Jolla, CA) in 1 ml of PBS rocking overnight at 4°C. The beads (referred to as LHR-IgG beads, CD4-IgG beads and huIgG-beads) were washed 3X in PBS and

35

once with lysis buffer. The CD4-IgG and huIgG beads were used as controls.

The precleared lysate described in Section A, above, was centrifuged at 10,000 x g for 10 sec, CaCl<sub>2</sub> was added to the supernatant at a final concentration of 5 mM, and the supernatant was mixed immediately with either LHR-IgG beads, CD4-IgG beads or huIgG-beads (typically 200 µl of precleared lysate per 10 µl packed beads), and incubated for 4 hr at 4°C on a rocker. The beads were washed 6X with lysis buffer, transferred to a new tube, and washed once more with lysis buffer.

The materials bound to the LHR-IgG beads were solubilized by boiling in SDS in the presence of 2-mercaptoethanol, electrophoresed on SDS-polyacrylamide gels (9 or 10%) and subjected to fluorography with ENTENSIFY or EN<sup>3</sup>HANCE (NEN). By fluorography, the 50 kd component tended to be more diffuse with ENTENSIFY than EN<sup>3</sup>HANCE. In the reprecipitation experiment, the SDS-solubilized sample was electrophoresed on a 7.5% SDS-gel with prestained standards (BioRad, high range) as markers. The region around 50 kd on the gel was excised by utilizing prestained ovalbumin (49.5 kd) as a position marker, and the protein electroeluted (BioRad model 422) into Laemmli running buffer at 60 mA overnight. The eluate was concentrated and the buffer was exchanged into 10 mM CHAPS in PBS on a Centricon 30 unit (Amicon, Danvers, MA), followed by incubation with LHR-IgG beads CD4-IgG or huIgG beads as described above. For the analysis of crude lysate, 200 µl of the precleared lysate was precipitated with cold acetone (80% v/v) and then subjected to electrophoresis as above.

LHR-IgG beads precipitated a diffuse 50 kd component (apparent molecular weight range is 50 kd - 58 kd) from [<sup>35</sup>S]-sulfate-labeled mesenteric lymph nodes (MLN) or peripheral lymph nodes (PN). A band of ≈90 kd (83 kd - 102 kd), relatively minor in terms of sulfate incorporation, was also observed in most analyses. In control precipitations, CD4-IgG and huIgG beads did not recognize the 50 kd major component or the 90 kd component in the lysates. When crude lysates were

directly analyzed, the 50 kd component represented the major constituent among several other bands. The tissue distribution of the 50 kd component was further examined by applying the identical protocol for [<sup>35</sup>S] sulfate-labeling and precipitation with LHR IgG to a number of organs. Among lymphoid tissues, only peripheral lymph nodes and mesenteric lymph nodes showed the 50 kd and 90 kd bands, while Peyer's patches, spleen, and thymus were negative for both. Non-lymphoid organs such as kidney, liver, cerebrum, and cerebellum were also completely negative.

LHR-IgG beads precipitated the 50 kd component when calcium was present, but not in its absence. The specificity of the interaction was further examined with the use of MEL-14 mAb. Preincubation of LHR-IgG beads with this antibody completely blocked the binding of the 50 kd band to the beads, whereas a class-matched control antibody (anti-CD45) had no effect. Fucoidin completely blocked the precipitation of the 50 kd component by LHR-IgG beads, while control polysaccharides (chondroitin sulfate B, chondroitin sulfate A, keratan sulfate) were completely inactive. Further, the presence of PPME significantly reduced the intensity of the 50 kd band, although a relatively high concentration was required. A control yeast mannan (mnn 2) had no effect at the same concentration. The precipitation of the minor 90 kd band by LHR-IgG beads was also calcium dependent, inhibitable by MEL-14 mAb, and blocked by fucoidin and PPME.

Finally, sialidase treatment of the glycoproteins was found to inhibit binding by LHR-IgG. Thus, sialic acid on the glycoproteins is apparently essential for binding. This result is in agreement with previous characterizations of interactions between selectins and their ligands.

The examples above demonstrate the ability of the assays of the present invention to identify effective competitive inhibitors of selectin-mediated intercellular adhesion. For the purposes of clarity and understanding, the invention has been described in these examples and the above disclosure in some detail. It will be apparent, however, that

certain changes and modifications may be practiced within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A method of assaying a test compound for the ability to inhibit intercellular adhesion mediated by a selectin receptor, the method comprising the steps of:

5       contacting the test compound with the receptor and an isolated receptor-binding agent, the agent being selected from the group consisting of a phosphorylated polysaccharide, a sulfated polysaccharide, a sulfated glycolipid, and a compound comprising the extracellular region of an endothelial cell  
10       surface glycoprotein; and

      detecting the ability of the test compound to inhibit binding between the receptor and the agent.

2. A method of claim 1 wherein the receptor is a  
15       lymphocyte homing receptor.

3. A method of claim 1 wherein the receptor is a chimeric molecule comprising a lymphocyte homing receptor component and an immunoglobulin component.  
20

4. A method of claim 1 wherein the receptor is immobilized on a solid surface.

5. A method of claim 5 wherein the step of detecting  
25       the binding of the receptor-binding agent to the receptor is carried out using flow cytometry.

6. A method of claim 1 wherein the receptor-binding agent is PPME.  
30

7. A method of claim 1 wherein the receptor-binding agent is fucoidin or dextran sulfate.

8. A method of claim 1 wherein the receptor binding  
35       agent is sulfatide.

9. A method of claim 1 wherein the glycoprotein is Sgp<sup>50</sup> or Sgp<sup>90</sup>.

10. A method of claim 1 wherein the receptor-binding agent is labeled.

11. A method of claim 10 wherein the receptor-binding agent is labeled with fluorescein.

12. A method of assaying a test compound for the ability to inhibit intercellular adhesion mediated by a lymphocyte homing receptor, the method comprising the steps of:  
10 contacting the test compound with the receptor immobilized on a solid surface and an isolated receptor-binding agent in a cell-free environment, the agent being selected from the group consisting of a phosphorylated polysaccharide, a sulfated polysaccharide, a sulfated glycoconjugate, and a  
15 compound comprising a vascular endothelial cell surface sulfated glycoprotein; and

detecting the ability of the test compound to inhibit binding between the receptor and the receptor-binding agent.

13. A method of claim 12 wherein the sulfated glycoprotein is Sgp<sup>50</sup> or Sgp<sup>80</sup>.

14. A method of claim 12 wherein the step of detecting inhibition of binding between the receptor and the  
25 receptor-binding agent is carried out using a first antibody directed against the receptor-binding agent and a second labeled antibody directed against the first antibody, whereby the receptor-binding agent is indirectly labeled.

15. A method of claim 12 wherein the receptor-binding agent is fluoresceinated and the step of detecting binding is carried out using a labeled antibody directed to fluorescein.

16. A method of assaying a test compound for the ability to inhibit intercellular adhesion mediated by a lymphocyte homing receptor, the method comprising the steps of:

in a cell-free environment, contacting the test compound with the receptor and a sulfated glycolipid capable of selectively binding the receptor, the glycolipid being immobilized on a solid surface; and

5 detecting the ability of the test compound to inhibit binding between the receptor and the glycolipid.

17. A method of claim 16 wherein the glycolipid is sulfatide.

10

18. A method of claim 16 wherein the receptor is a chimeric molecule comprising a lymphocyte homing receptor component and an immunoglobulin component.

15

19. A method of assaying a carbohydrate test compound for the ability to inhibit intercellular adhesion mediated by a lymphocyte homing receptor, the method comprising the steps of:

20 contacting the test compound with a fluoresceinated receptor-binding agent and a lymphocyte; and

detecting the ability of the test compound to inhibit binding of the receptor-binding agent to the cell using a fluorescence activated flow cytometer.

25

20. A method of claim 19 wherein the fluoresceinated receptor-binding agent is selected from the group consisting of a sulfated polysaccharide, a sulfated glycolipid, and a compound comprising the extracellular region of an endothelial cell surface glycoprotein

30

21. A method of claim 19 wherein the fluoresceinated receptor-binding agent is fucoidin.

22. A composition comprising an isolated inhibitor of selectin-mediated intercellular adhesion identified by a method comprising the steps of:

35

contacting the inhibitor with a selectin receptor and an isolated receptor-binding agent, the agent being selected

from the group consisting of a phosphorylated polysaccharide, a sulfated polysaccharide, a sulfated glycolipid, and a compound comprising the extracellular region of an endothelial cell surface glycoprotein; and

5 detecting the ability of the inhibitor to inhibit binding between the receptor and the agent.

23. A composition of claim 22 wherein the selectin receptor is a lymphocyte homing receptor.

10 24. A composition of claim 22 wherein the selectin receptor is immobilized on a solid surface.

15 25. A composition of claim 22 wherein the receptor-binding agent is fluoresceinated.

26. A composition of claim 22 wherein the receptor-binding agent is PPME, fucoidin, dextran sulfate, sulfatide, Sgp<sup>50</sup> or Sgp<sup>90</sup>.

20 27. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an inhibitor in accordance with claim 22.

25



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/03606

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12Q 1/00; G01N 33/53

US CL : 435/7.24, 7.1, 7.2, 7.92

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.24, 7.1, 7.2, 7.92

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG, BIOSIS

search terms: selectin receptor, adhesion, inhibitor

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Journal of Cell Biology, Volume 104, issued March, 1987, T.A. Yednock et al., "Phosphomannosyl-derivatized Beads Detect a Receptor Involved in Lymphocyte Homing", pages 713-723, see page 714.	1-27
Y	Journal of Cell Biology, Volume 111, issued September, 1990, Y. Imai et al., "Direct Demonstration of the Lectin Activity of gp90MEL, a Lymphocyte Homing Receptor", pages 1225-1232, see whole article.	1-27
Y	Journal of Cell Biology, Volume 104, issued March, 1987, T.A. Yednock et al., "Receptors Involved in Lymphocyte Homing: Relationship Between a Carbohydrate-binding Receptor and the MEL-14 Antigen", pages 725-731, see whole article.	1-27

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

29 JULY 1992

Date of mailing of the international search report

06 AUG 1992

Name and mailing address of the ISA/  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

DAVID GUZO

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-1906